

Shot record
depth
migration of
georadar

R. J.
Ferguson, A.
Pidlisecky,
and C. Rowell

Outline

Introduction

Data
preparation

Data
processing

Imaging

Conclusions

Future work

Acknowledgements

Shot record depth migration of georadar

R. J. Ferguson, A. Pidlisecky, and C. Rowell



Shot record
depth
migration of
georadar

R. J.
Ferguson, A.
Pidlisecky,
and C. Rowell

Outline

Introduction

Data
preparation

Data
processing

Imaging

Conclusions

Future work

Acknowledgements

1 Introduction

2 Data preparation

3 Data processing

4 Imaging

5 Conclusions

6 Future work

7 Acknowledgements

Introduction

- Low dielectric permittivity in basalt causes strong electromagnetic reflections.
- Low conductivity of basalt allows a great depth of penetration.
- A large basalt flood at Craters of the Moon, Idaho contains a plumbing system of volcanic conduits.
- Conduits are metres in height and circumference — need sub-decimetre trace spacings.

Shot record
depth
migration of
georadar

R. J.
Ferguson, A.
Pidlisecky,
and C. Rowell

Outline

Introduction

Data
preparation

Data
processing

Imaging

Conclusions


Future work

Acknowledgements


Data preparation

- Survey normalization and rotation.
- Survey regularization.
- Trace regularization.

Survey



Survey regularization



Shot record
depth
migration of
georadar

R. J.
Ferguson, A.
Pidlisecky,
and C. Rowell

Outline
Introduction

**Data
preparation**

Data
processing



Imaging

Conclusions

Future work

Acknowledgements

Trace regularization



Shot record
depth
migration of
georadar

R. J.
Ferguson, A.
Pidlisecky,
and C. Rowell

Outline

Introduction

Data
preparation

Data
processing

Imaging

Conclusions

Future work

Acknowledgements

Data processing

- Determine frequency range.
- Spiking deconvolution.
- Gabor deconvolution.

Shot record
depth
migration of
georadar

R. J.
Ferguson, A.
Pidlisecky,
and C. Rowell

Outline

Introduction

Data
preparation

Data
processing


Imaging

Conclusions

Future work

Acknowledgements

Raw data



Shot record
depth
migration of
georadar

R. J.
Ferguson, A.
Pidlisecky,
and C. Rowell

Outline

Introduction

Data
preparation

Data
processing


Imaging

Conclusions

Future work

Acknowledgements

Spiking decon



Shot record
depth
migration of
georadar

R. J.
Ferguson, A.
Pidlisecky,
and C. Rowell

Outline

Introduction

Data
preparation

Data
processing


Imaging

Conclusions

Future work

Acknowledgements

Gabor decon



Shot record
depth
migration of
georadar

R. J.
Ferguson, A.
Pidlisecky,
and C. Rowell

Outline

Introduction

Data
preparation

Data
processing

Imaging

Conclusions

Future work

Acknowledgements

Imaging

- ZOM vs. PSDM.
- ZOM: Gabor + elevation statics → ZOM.
- PSDM: Gabor → PSDM.

Shot record
depth
migration of
georadar

R. J.
Ferguson, A.
Pidlisecky,
and C. Rowell

Outline

Introduction

Data
preparation

Data
processing




Imaging

Conclusions

Future work

Acknowledgements

Migration



Shot record
depth
migration of
georadar

R. J.
Ferguson, A.
Pidlisecky,
and C. Rowell

Outline

Introduction

Data
preparation

Data
processing



Imaging

Conclusions

Future work

Acknowledgements

Zoom



Shot record
depth
migration of
georadar

R. J.
Ferguson, A.
Pidlisecky,
and C. Rowell

Outline
Introduction

Data
preparation

Data
processing


Imaging

Conclusions

Future work


Acknowledgements

Beauty cave



From Rowell et al, 2010, *Characterization of lava tubes using ground penetrating radar at Craters of the Moon National Monument, USA*: CREWES Research Report.

Stratified flows



From Rowell et al, 2010, *Characterization of lava tubes using ground penetrating radar at Craters of the Moon National Monument, USA*: CREWES Research Report.

Conclusions

- Sub-decimetre georadar acquisition is migrateable.
- Deconvolution works: Gabor deconvolution pulls out deep reflections.
- PSDM is superior to elevation statics + ZOM.

Shot record
depth
migration of
georadar

R. J.
Ferguson, A.
Pidlisecky,
and C. Rowell

Future work

- Velocity analysis.
- Longer traces.
- Tighter line spacing.
- Real 3D over an interesting target.

Shot record
depth
migration of
georadar

R. J.
Ferguson, A.
Pidlisecky,
and C. Rowell

Outline

Introduction

Data
preparation

Data
processing

Imaging

Conclusions

Future work

Acknowledgements

Acknowledgements

- CREWES sponsors and NSERC.
- Dr. James D. Irving, School of Engineering, University of Guelph.
- Kevin W. Hall.